



HEAD INJURY AND CONCUSSION POLICY

Introduction

Langley School recognises that all concussions or suspected concussions must be taken seriously to safeguard the safety and long-term health of those affected.

Everyone has a role to ensure that they are appropriately informed and understand what role they play in prevention, recognition, and management of suspected concussion.

Concussion is not always a visible injury, and in particular, children and adolescents can take longer to recover because their brains are still developing, therefore, a more conservative approach should be taken.

Policy Aim

The aim of this policy is to:

- Ensure understanding of the key terms and the link between head injury and brain injury.
- Identify sport activities which carry a risk of head injury.
- Underscore the importance of creating suitable risk assessments for sport activities being undertaken by the school; and
- Provide clear processes to follow when a student does sustain a head injury.

This policy applies to:

- School staff (including part time or occasional employees or visiting teachers);
- Pupils/students of the School
- Parents of pupils/students at the school; and
- Any other individual participating in any capacity in a school activity. For example, this would include a contractor providing sports coaching, or a volunteer on a school trip.

A head injury could happen in any area of School life. This policy focuses on sport activities (both contact sports and non-contact sports) where the risk of head injuries happening is higher but can be used for head injuries which occur in another context.

Definitions

The following terms are used in this policy:

Head injury: means any trauma to the head other than superficial injuries to the face.

Traumatic Brain Injury (TBI): is an injury to the brain caused by a trauma to the head (head injury).

Concussion: is a type of traumatic brain injury (TBI) resulting in a disturbance of brain function. It usually follows a blow directly to the head, or indirectly if the head is shaken when the body is struck. Transient loss of consciousness is not a requirement for diagnosing concussion and occurs in less than 10% of concussions.

Transient Loss of consciousness is the sudden onset, complete loss of consciousness of brief duration with relatively rapid and complete recovery. It can also be referred to as 'being knocked out' or a 'blackout.'

Persistent loss of consciousness is a state of depressed consciousness where a person is unresponsive to the outside world. It can also be referred to as a coma.

Chronic Traumatic Encephalopathy (CTE) is one type of degenerative and progressive brain condition that's thought to be caused by TBIs and repeated episodes of concussion. CTE usually begins gradually several years after receiving TBIs or repeated concussions. The symptoms affect the functioning of the brain and eventually lead to dementia.

Contact sport: is any sport where physical contact is an acceptable part of play for example rugby, football, and hockey.

Non-contact sport is any sport where physical contact is not an acceptable part of play but where there are nonetheless potential collisions between players and between players and the ball, for example cricket and netball.

The risks

- Playing contact and non-contact sport increases an individual's risk of collision with objects or other players.
- Collisions can cause a head injury, which can cause a traumatic brain injury such as a concussion.
- It is very important to recognise that a pupil/student can have a concussion, even if they are not 'knocked out'. Transient loss of consciousness is not a requirement for diagnosing concussion and occurs in less than 10% of concussions.
- Children and young adults are more susceptible to concussion than adults because their brains are not yet fully developed and thus more vulnerable to injury.
- The current evidence suggests that repeated episodes of concussion, even where there is no transitory loss of consciousness, can cause significant changes to the structure and function of the brain in a condition known as Chronic Traumatic Encephalopathy (CTE).

Raising Awareness

- Head injury awareness will be regularly revisited with pupils/student (age appropriate) this may include focused talks and/or information posters.
- This policy will be available to parents and staff on the school website and via the staff launch page, under policies, whole school.
- The RFU Headcase resources are available to all via [07. HEADCASE \(keepyourbootson.co.uk\)](#) training can be accessed free via [Courses \(keepyourbootson.co.uk\)](#) selecting the 'book a course' tab and filtering to 'virtual courses'. All staff delivering sports and our school medical team will be required to undertake the course.

Policy author: Head of Sport/Director of Compliance and Operations

Date: Lent 2023

Review date: Lent 2024

Preventative steps to reduce the risks

Any person responsible for the undertaking of a sporting activity must ensure a suitable risk assessment for the specific sport activity is created.

This risk assessment should be tailored to the specific School environment and should:

- Identify the specific risks posed by the sport activity, including the risk of players sustaining head injuries.
- Identify the level of risk posed.
- State the measures and reasonable steps taken to reduce the risks and;
- Identify the level of risk posed with the measures applied.

The governing bodies of most sports played in Schools have each produced head injury guidelines that are specific to their sport. Those responsible for risk assessing sport activities in School should have regard to the relevant and latest guidelines when carrying out their risk assessment. For example:

The Sport and Recreation Alliance includes members from the major sports governing bodies, including the RFU, ECB, FE, RFL and England Hockey. Together they have produced ‘Concussion Guidelines for the Education Sector’, which can be viewed here: https://www.afpe.org.uk/physical-education/wp-content/uploads/Concussion_guidelines_for_the_education_sector_June2015.pdf

Football

General FA concussion guidelines: <https://www.thefa.com/get-involved/fa-concussion-guidelines-if-in-doubt-sit-them-outold>

FA Heading Guidance: <https://www.thefa.com/news/2020/feb/24/updated-heading-guidance-announcement-240220>

Rugby

<https://www.englandrugby.com/participation/playing/headcase/age-grade/schools-and-colleges>

RFU Graduated Return to Play guidelines: <https://www.englandrugby.com/dxdam/04/0453acb5-5fe2-4608-91b0-a2bd191c3016/HEADCASE%20GRTP.pdf>

Hockey

GB & England Hockey Concussion Policy <https://www.cuhc.org.uk/wp-content/uploads/2020/10/CUHC-Concussion-Policy-2020-21.pdf>

England Hockey ‘Safe Hockey’ guides <https://www.englandhockey.co.uk/governance/duty-of-care-in-hockey/safe-hockey>

Policy author: Head of Sport/Director of Compliance and Operations

Date: Lent 2023

Review date: Lent 2024

Potential measures to reduce the risk of players sustaining head injuries while playing sports might include:

- Structuring training and matches in accordance with current guidelines from the governing body of the relevant sport (see above).
- Removing or reducing contact elements from contact sports, for example removing 'heading' from football.
- Removing or reducing the contact elements of contact sports during training sessions.
- Ensuring that there is an adequate ratio of coaches to players in training.
- Ensuring that students are taught safe playing techniques.
- Ensuring that students are taught to display sportsman like conduct at all times and maintain respect for both opponents and fellow team members equally.
- Using equipment and technology to reduce the level of impact from collision with physical objects (e.g., using padding around rugby posts, using soft balls, not overinflating footballs etc.).
- Using equipment and technology to reduce the level of impact from collision between players (e.g., gumshields, helmets etc).
- Coaching good technique in high-risk situations (such as rugby tackles).
- Ensuring that the playing and training area is safe (for example, that is not frozen hard, and there are suitable run-off areas at the touchlines).
- Ensuring that a medical professional is easily accessible during training and matches.

Head injuries sustained outside of school

As noted above, repeated concussions can cause significant changes to the structure and function of the brain, in particular the child's brain.

It is therefore very important that Langley school, pupils, students, and parents take a holistic approach to the management of head injury causing concussions and cooperate with regards to sharing information.

Where a student sustains a head injury which has caused a concussion whilst participating in an activity outside of school, parents of the student concerned should promptly provide the school nurses with sufficient details of the incident, and keep the school updated of any developments thereafter. This would apply, for example, if a student suffers a concussion playing rugby for an external rugby club or if a student sustains a head injury while taking part in an informal game of sport, for example in the local park.

The school will determine the appropriate way forward on receiving a notification of this nature. That might include reviewing any return to play plan already established by the external club, or if no such plan has been put in place, considering whether a return to play plan should be established under this policy.

In turn the school will inform parents where a student has sustained a head injury causing a concussion at School.

Policy author: Head of Sport/Director of Compliance and Operations

Date: Lent 2023

Review date: Lent 2024

Procedure to follow where a student sustains a head injury at School

The welfare of students is of central importance. Any person to whom this policy applies should adopt a cautious approach if they are in any doubt as to whether a head injury has occurred and/or whether the head injury has caused a concussion.

Where a student sustains a suspected head injury or concussion, the person supervising the activity should immediately remove the student from play where it is safe to do and seek appropriate medical professional from the school nurse or a qualified first aider.

Those individuals to whom this policy applies should be aware of the symptoms of a concussion. The British Medical Journal has published a one page 'Pocket Concussion Recognition Tool' to help identify concussion in children, youth and adults. The tool is attached at Schedule Two, and is also available for download (here: <https://bjsm.bmj.com/content/bjsports/47/5/267.full.pdf>) The tool identifies the following signs and symptoms of suspected concussion:

- Loss of consciousness
- Seizure or convulsion
- Balance problems
- Nausea or vomiting
- Drowsiness
- More emotional
- Irritability
- Sadness
- Fatigue or low energy
- Nervous or anxious
- "don't feel right"
- Difficulty remembering
- Headache
- Dizziness
- Confusion
- Feeling slowed down
- "Pressure in head"
- Blurred vision.
- Sensitivity to light.
- Amnesia.
- Feeling like "in a fog"
- Neck pain
- Sensitivity to noise; and
- Difficulty concentrating

Where a pupil/student display any of the symptoms above, they should not be permitted to return to play and should be assessed by the medical professional.

Policy author: Head of Sport/Director of Compliance and Operations

Date: Lent 2023

Review date: Lent 2024

The medical professional should determine whether the student is displaying any “red flag” symptom in which case the ambulance services should be called on 999. The Pocket Concussion Recognition Tool at Appendix ii, below, identifies the following red flags:

- Athlete complains of neck pain
- Increasing confusion or irritability
- Repeated vomiting
- Seizure or convulsion
- Weakness or tingling/burning in arms or legs
- Deteriorating conscious state
- Severe or increasing headache
- Unusual behaviour change; and
- Double vision.

The school must ensure that the pupil/student’s parents are notified of the head injury as soon as reasonably possible, and in any case on the same day of the incident, please see individual setting procedures a ‘**Advice to parents/carers of children following a head injury**’, below. Please ensure you note this has been provided on the AccidentBook entry (as below).

Anyone sustaining a head injury and showing symptoms of concussion will not be allowed to drive themselves or travel home unaccompanied by either school or public transport.

Whenever a pupil/student suffers a suspected head injury the attending member of staff must report this via the schools AccidentBook system, which can be accessed from the school’s staff launch page, please also ensure that the form is updated if advice and treatment changes.

Managing a return to play following a head injury

Any student that has suffered a head injury and showed symptoms of concussion should be subject to a graduated return to play programme (**GRTP**).

The GRTP should be developed in consultation with a suitably qualified sporting/medical professional and be tailored to the specific circumstances of the individual (including the type of injury sustained and the relevant sport). For an example GRTP, see the GRTP developed by

England Rugby here: <https://www.englandrugby.com/dxdam/04/0453acb5-5fe2-4608-91b0-a2bd191c3016/HEADCASE%20GRTP.pdf>

Staff must ensure GRTP information is transferred to the off games sheets, where applicable.

It is the responsibility of the parents to ensure that their child does not participate in any inappropriate physical activity outside of school whilst they are subject to a GRTP.

If the child participates in school-sponsored sports within this period, they must have a letter from a Licensed Health Care Provider (LHCP) clearing them for return to play eligibility. LHCPs are defined as: Medical Doctors (MD), Doctors of Osteopathy (DO), Advanced Registered Nurse Practitioners (ARNP), Physician Assistants (PA-C), or Certified Athletic

Policy author: Head of Sport/Director of Compliance and Operations

Date: Lent 2023

Review date: Lent 2024

Trainers (AT/L). If you seek medical care, please provide the school with documentation from a Licensed Health Care Provider for any activity limitations or restrictions.

Breaches of this policy

The school takes its duty of care very seriously. The school will take appropriate action against any person found to have breached this policy. For example:

- if a student attempts to return to play in breach of their GRTP plan, the school would consider the matter under the school's behaviour policy;
- if a member of staff fails to report a head injury, the school would consider the matter under the school's staff disciplinary policy; and
- if a parent fails to report to the school a head injury their child sustains outside of school, the school will consider the matter under the terms of the school parent contract.

Advice to parents/carers of children following a head injury

Note: Usual parent contact procedures should be followed, in accordance to setting. In addition, the following information should be provided to parent/carer.

Pre-Prep/Nursery – Please ensure the accident report head injury box is ticked and that the form includes head injury information. This should be given to the parent on the day of the injury.

Prep – Please ensure parents are provided with the Prep School Head Injury Advice Sheet on the day of the injury.

Senior – Please ensure parents are provided with the Senior School Head Injury Advice Sheet on the day of the injury.

Due to the inconsistent nature of head injuries, children who have received even what is seemingly a slight bump on the head should be observed for at least 24 hours after the accident. Symptoms of a head injury can be delayed for several hours or even a day following the injury.

Policy author: Head of Sport/Director of Compliance and Operations

Date: Lent 2023

Review date: Lent 2024

Appendix ii Concussion Recognition Tool

Pocket CONCUSSION RECOGNITION TOOL™

To help identify concussion in children, youth and adults



RECOGNIZE & REMOVE

Concussion should be suspected **if one or more** of the following visible clues, signs, symptoms or errors in memory questions are present.

Annexure 1 Visible clues of suspected concussion

Any one or more of the following visual clues can indicate a possible concussion:

- Loss of consciousness or responsiveness
- Lying motionless on ground / Slow to get up
- Unsteady on feet / Balance problems or falling over / Incoordination
- Grabbing/Clutching of head
- Dazed, blank or vacant look
- Confused / Not aware of plays or events

Annexure 2 Signs and symptoms of suspected concussion

Presence of any one or more of the following signs & symptoms may suggest a concussion:

- Loss of consciousness
- Seizure or convulsion
- Balance problems
- Nausea or vomiting
- Drowsiness
- More emotional
- Irritability
- Sadness
- Fatigue or low energy
- Nervous or anxious
- "Don't feel right"
- Difficulty remembering
- Headache
- Dizziness
- Confusion
- Feeling slowed down
- "Pressure in head"
- Blurred vision
- Sensitivity to light
- Amnesia
- Feeling like "in a fog"
- Neck Pain
- Sensitivity to noise
- Difficulty concentrating

© 2013 Concussion in Sport Group

3. Memory function

Failure to answer any of these questions correctly may suggest a concussion.

- "What venue are we at today?"
- "Which half is it now?"
- "Who scored last in this game?"
- "What team did you play last week / game?"
- "Did your team win the last game?"

Any athlete with a suspected concussion should be IMMEDIATELY REMOVED FROM PLAY, and should not be returned to activity until they are assessed medically. Athletes with a suspected concussion should not be left alone and should not drive a motor vehicle.

It is recommended that, in all cases of suspected concussion, the player is referred to a medical professional for diagnosis and guidance as well as return to play decisions, even if the symptoms resolve.

RED FLAGS

If ANY of the following are reported then the player should be safely and immediately removed from the field. If no qualified medical professional is available, consider transporting by ambulance for urgent medical assessment:

- Athlete complains of neck pain
 - Increasing confusion or irritability
 - Repeated vomiting
 - Seizure or convulsion
 - Deteriorating conscious state
 - Severe or increasing headache
 - Unusual behaviour change
 - Double vision -
- Weakness or tingling / burning in arms or legs

Remember:

- In all cases, the basic principles of first aid (danger, response, airway, breathing, circulation) should be followed.
- Do not attempt to move the player (other than required for airway support) unless trained to do so
- Do not remove helmet (if present) unless trained to do so.

from McCrory et. al, Consensus Statement on Concussion in Sport. Br J Sports Med 47 (5), 2013

© 2013 Concussion in Sport Group

Policy author: Head of Sport/Director of Compliance and Operations

Date: Lent 2023

Review date: Lent 2024